

An integrated approach on satellite geodesy data to delineate morphotectonic and neotectonic activities in the West Coast off Kerala, Southern India

Drishya Girishbai¹ and Arun Bhadran²

Received 7 October 2017; accepted 9 February 2018; published 19 March 2018.

Better understanding of marine geomorphology plays a vital role in the development of humankind. Therefore, satellite altimetry data serves as a comprehensive and cohesive method. The present study attempts to explore the delineation of submarine structural features and a correlation of tectonic features both on land and offshore of the west coast off Kerala, using Global bathymetric model (GBM) and satellite-derived gravity data. The study observes the south-west coast of Indian peninsula, which has a unique geological and structural features. It is a passive margin, which has undergone drift-rift mechanism associated with the separation of Madagascar–Seychelles–India. The hinterland of the area is an assortment from the Meso-Archean to Neo-Proterozoic crustal blocks sandwiched with shear zones. The study area has been divided into three zones (Zone I, II and III) based on major shear zones in continental part to decipher the tectonic control. Satellite altimetry data has been used to derive drainages with different threshold to study submarine drainage system, channel shift, fault, impact crater etc. Conversely, all these observations deduced by satellite altimetry data have been validated with available published data. This study implies that geodesy data is an important and economical tool for preliminary studies, for understanding regional geological niche and its structural/tectonic control.

KEYWORDS: Satellite altimetry; neotectonism; passive margin; oblique rift valley.

Citation: Girishbai, Drishya and Arun Bhadran (2018), An integrated approach on satellite geodesy data to delineate morphotectonic and neotectonic activities in the West Coast off Kerala, Southern India, *Russ. J. Earth. Sci.*, 18, ES2003, doi:10.2205/2018ES000617.

1. Introduction

Scientific ocean exploration started since Her Majesty's Ship Beagle, and nowadays voyages, us-

¹International Affairs & IGC, Central Head Quarters, Geological Survey of India, Kolkata, India

²Geodata Division, Central Head Quarters, Geological Survey of India, Kolkata, India

Copyright 2018 by the Geophysical Center RAS.
<http://rjes.wdcb.ru/doi/2018ES000617-res.html>

ing high resolution seismic and sequence stratigraphic studies continue to make discoveries. These studies are essential for sustainable future of humankind and even interplanetary voyages. However, the expenditure towards such studies are enormous and economical probes are essential for disabling those difficulties. The satellite altimetry and remote sensing is an alternate method for the preliminary investigation of marine geomorphological characterization in the regional scale. It also provides the elevation frequency distribution (hypsometry) in the continental and oceanic area which

can decipher large-scale manifestations of plate tectonic features [*Gahagan et al., 1988; Sandwell and Smith, 1994*].

More than half of the Earth's continental and continental shelf part is covered with sediments and sedimentary rocks, with maximum accumulations at continental rift margins. Submarine channel-levee complexes act as a conduit for terrigenous material from continental margins to the deep oceans. Details on the evolutionary processes of channel play a major role for the predictions of their individual occurrence, morphology, geohazard zone etc.; which are essential for seafloor infrastructure planning and construction [*Allen, 2008; Paull et al., 2003; Xu, 2010*].

Indian peninsula has a unique feature: it is bounded by world's largest sedimentary basins like Bengal fan on east and Indus fan on west. Ocean basin calculated by satellite altimetry data has been widely used in Indian Continent mainly for delineating major drainage networks in submarine fans of Bengal, Indus and Ayyeyarwady in Andaman Sea [*Girishbai et al., 2017; Kundu and Pattnaik, 2011; Prerna Ramesh et al., 2015*]. In this paper a detailed investigation has been made to delineate submarine drainages, morphotectonic and neotectonic activities in the west coast off Kerala using Global bathymetric model (GBM) and satellite derived gravity data with an aid of Geographic Information System (GIS) software.

2. Study Area and Regional Geology

The linear feature of the West coast of Indian peninsular (i.e., from Sir Creek in the north to Kanyakumari in the south) is of drift-rift tectonism of Indian Plate since 167 Ma (i.e., after the breakup of eastern Gondwana) that cause the separation of India, Madagascar and Seychelles (~ 90 ma and ~ 65 ma) [*Chatterjee et al., 2013; Norton and Sclater, 1979*] along with magmatic plume activity. In the entirety west coast of India act as a passive margin and accelerate the voyage and collision of Indian Plate with the Eurasian plate (Himalayan Orogeny), which makes west coast of India, a unique place to explore tectonic features.

The present study area is bounded by Precambrian basement rocks in the east known as Southern Granulite Terrain (SGT), which are followed

by a hiatus and overlies by Mesozoic dykes and Cenozoic sediments. Toward the west the continental shelf forms the Kerala-Konkan (KK) Basin, which is the sub-basin of Laccadive Basin (LB). LB dissected by Vishnu Fracture Zone (VFZ), the Laccadive-Chagos Ridges (LCR) [*Bhattacharya and Yatheesh, 2015; Gopala Rao et al., 2010*], which occur on the west of the basin and act as a vital part of oblique rift valley [*Subrahmanyam and Chand, 2006*].

The Basement rocks i.e., SGT, is an amalgamation of Mesoproterozoic to Neoproterozoic crustal blocks exposing mid-lower crustal rocks dissected by Palghat-Cauvery Shear System (includes Moyar-Bhavani, Palghat-Cauvery and Attur shear zones) and Achankovil Shear Zone (ACSZ). These shear zones divide the southern part of SGT from north to south into three major crustal blocks known as Coorg, Madurai and Trivandrum [*Chetty et al., 2016; Collins et al., 2014; Santosh et al., 2009*] (Figure 1). The marginal part of the west coast of India, is mostly controlled by three principal structural trends (Dharwar (NNW-SSE), Aravalli orogenies (NE-SW) and Narmada graben (E-W), which presents a straight coastline, with the continental shelf width varying from 345 km off Daman in the north to 120 km off Goa and tapering to 60 km off Kochi to the south [*Faruque and Ramachandran, 2014*].

During the Late Mesozoic the region experienced thermal regime associated with the breakup of Gondwana resulted pericratonic continental fracturing and emplacement of basaltic dykes in three major episodes (~ 145 Ma, $\sim 90 - 85$ Ma and $\sim 70 - 65$ Ma), which made different stage of west coast evolution [*Ajayakumar et al., 2017; Karunakaran and Mahadevan, 1971; Radhakrishna, 1999*]. The KK Basin, located on the continental shelf part has evolved by separation of continental masses from western margin of India and suggest it as an oceanic crust between LCR and KK shelf [*Das, 2013*]. LCR is the most significant topographic feature with several islands observed along its strike length. This ridge is recognized as a hot spot trace related to the Reunion plume [*Duncan and Storey, 1992*]. Furthermore, Kerala deep-water basin, is bounded by two prominent north-south oriented fracture zones of the Indian Ocean, known as Vishnu fracture zone (in the west) and Indrani fracture zone (in the east). VFZ, a trans-




Figure 1. Generalized tectonic framework of Southern India illustrating major crustal blocks and intervening shear/suture zones (after [Collins et al., 2014; Santosh et al., 2015, 2017]).

form fault which cause the separation of Seychelles from India and it extends from the Kerala shelf to the Carlsberg-Ridge, over a length of more than 2500 km [Bastia et al., 2010; Bhattacharya and Yatheesh, 2015] (Figure 2). In addition to all tectonic features mentioned above, the Geoidal gravitational low observed along the study area makes it a unique place.

3. Material and Methods

GIS-based hypsometric data is used extensively on land for detecting streams and channels [Belknap and Naiman, 1998; Imaki and Beechie, 2008].

It became an integral part of hydrologic studies due to the spatial character and its control on the hydrological processes [Vieux, 2001]. In the current study, an effort has been made to find the credibility of GIS on deep sea channels in the Arabian Sea along with the global seafloor topography datasets. To scrupulously constrain the submarine drainage patterns, data from http://topex.ucsd.edu/cgi-bin/get_data.cgi has been used, that pertains to v16.1 of the GBM data compiled by [Sandwell and Smith, 1997]. Data was downloaded in .xyz format and converted into a Digital Bathymetric Model (DBM). Further, Gravity data (mGal) has been used for the validation of topographic data. It was generated at a grid resolution of 1.7×1.7 km. High resolution elevation data were collected from Li-




Figure 2. (a) – Satellite gravity of the western Indian Ocean showing the various tectonic elements referred in the present study. (Modified after *Sandwell and Smith* [2009] and *Ramana et al.* [2015]). (b) – DEM of West Coast of Kerala (from Topex data) with major Shear Zones (McSZ: Mercara Shear Zone, MSZ: Moyar Shear Zone, PCSZ: Palaghat–Cauvery Shear Zone, KKPT: Karur–Kamabam–Painavu–Trichur, ACSZ: Achankovil Shear Zone) and major submarine features like Laccadivine-Chagos Ridge (LCR) Laccadive Basin, Allapey-Trivandrum Terrace Complex (ATTC), Pratap Ridge, Vishnu fracture Zone (VFZ) (Modified after [*Collins et al.*, 2014; *Santosh et al.*, 2015, 2017; *Yatheesh et al.* [2013]).

Table 1. Drainage extracted using different threshold value and its ordering

Sl No	Stream Threshold Value	Total Streams	# Land (Unclassified) Channels	# Ocean Channels	1st Order	2nd Order	3rd Order	4th Order
1	500	35	2	33	31	2	-	-
2	250	151	28	123	107	16	-	-
3	100	433	74	359	298	55	6	-
4	50	900	148	752	584	149	18	1

DAR, SONAR etc., which are specifically suited for managing, processing and integration of three dimensional data (Childs, C., 2011, Terrain Datasets: The Top 10 Reasons to Use Them. Arc User Online, Spring Edition).

4. Result and Discussions

Availability of satellite-derived high resolution Digital Elevation model (DEM) and spatial information technologies has gained rapid attention especially on morphotectonic analysis using geomorphic indices [Ferraris *et al.*, 2012; Reinfeld *et al.*, 2004]. DEM is the best digital tool to automate the morphological characterisation of drainages, their area of influence and hypsometric parameters [Rahman *et al.*, 2010]. O'Callaghan and Mark [1984] introduced the computation for DEM pixels based on flow routing model referred as D8 algorithm. Later, Jensen and Domigree [1988] has modified it by omitting the sink/depression. It is noted that omission of depressions/sink have a positive influence on the enhancement of the quality of elevation models [Lin *et al.*, 2008]. These sinks may exist as single pixels or alternatively, as a group of multiple pixels with values lesser than all the pixels surrounding them. The process to fill up these sinks may be direct or iterative in nature depending on the number of pixels involved. The DBM generated for the study area, has been corrected for erroneous depressions.

Binary raster network denoting drainage has been deduced from flow accumulation and ordering of drainages has been implemented by methods of [Horton, 1945; Shreve, 1966; Strahler, 1952]. After a cogitation of the channels within their network, basins were also delineated based on channel

flow direction and order. Figure 3 and Table 1 are created by adding different threshold values (500, 250, 100 and 50) for the better understanding of the drainage developments. Major sediment distribution along the west coast off Kerala is ranging from terrigenous to carbonaceous sand and clay [Rao and Wagle, 1997] (Figure 4). Based on drainage pattern and structural control three zones are selected for detailed study (Figure 5).

4.1. Zone I

In Zone I, the continental area is bound by Mercara and Bhavani Shear Zones. The Continental shelf is wider and more open towards north with minor protuberances of positive heights at 1500 isobaths, which is a part of Pratap Ridge, a coast parallel running ridge (Figure 5 and profile A-A' of Figure 6). The submarine drainages are parallel and their Hortonian higher order [Horton, 1945] shows a tilting/sinuosity in the NNW and then towards the SSE. The sediment distribution in the inner shelf of Zone I is mainly of modern clayey silt and silty clay with high organic matter (low carbonate content) and with less heavy minerals. On the other hand, mid-shelf is uneven and outer shelf is commonly interrupted by shore-parallel ridges and reefs [Faruque and Ramachandran, 2014; Rao and Wagle, 1997]. Similar morphotectonic studies on land has been carried out by [Vijith *et al.*, 2015] in Mahe River, Northern Kerala. They observed that tilt in river mouth (in NNW and SSE) from head water, implies neotectonic activities. So the sinuosity observed on submarine drainage may be of shear zone extension. Furthermore, this observation has been exemplified by the magnetic studies of [Gopala Rao *et al.*, 2010; Subrahmanyam and Chand, 2006; Subrahmanyam *et al.*, 1993] on the




Figure 3. Drainage with thresholds values: a – 500, b – 250, c – 100, and d – 50 values. *Editorial note:* To zoom selected image click on the corresponding number in red square, make one more click to return back.




Figure 4. Sedimentary distribution of west coast of Kerala and exhibit classification of Zones I–III based on Major Shear Zones on land.

extension of Palaghat Cauvery Lineament (mega lineament).

4.2. Zone II

Central part of the study area, i.e., Zone II, belongs to Madhurai Block which is bounded by E–W running Palaghat Cauvery shear zone (PCSZ) in

the North and South by Achankovil Shear Zone (ACSZ) of Pan African Orogeny with the NW–SE trend. Moreover, the continental part of the area includes the Western Ghats major break, known as Palaghat Gap along with the most active Karur–Kamabam–Painavu–Trichur (KKPT) lineament [Rajendran and Rajendran, 1996; Singh, 2016]. Satellite altimetry data has given signatures of the mount like features at 2000 m isobath; with




Figure 5. Major Features identified using topex/GBM data; features like submarine palaeochannels and impact craters are kept as inset.

submarine dendritic drainage pattern and a deep cut valley off Ponnani. *Faruque and Ramachandran* [2014] revealed that these deep cut valleys are the structural continuity of Palaghat Gap. Two major submerged paleo channels observed south off Ponnani. These paleo channels are supported by prominent occurrence of sand [*Rao et al., 2003; Sukumaran et al., 2010*] and existence of paleo channels [*Hashimi et al., 1995; Shareef et al., 2015*]

(Figure 4 and Figure 5). Furthermore, most rivers in the central part of Kerala show highly sinuous nature and asymmetric basin characteristics which indirectly point towards the planform changes of the river and watershed as a whole in response to the neotectonic activities in KKPT. Recent studies indicate that, minor tremors are occurring with KKPT, suggesting a significant tectonic activity [*Ajayakumar et al., 2017; Dhanya and Renoy, 2015;*




Figure 6. Maps with profiles showing major tectonic and morphological features in Zones I–III (A–A', B–B', C–C', and D–D').

Girish Gopinath et al., 2016; Rajendran and Rajendran, 1996].

An isolated circular depression observed south off Kochi (Figure 5 and profile B–B' of Figure 6).

Pillay et al. [2016] marked this feature as an impact crater. Moreover, coast parallel isobaths in Zone I (NNW–SSE) show a sudden shift in Zone II (NNE–SSW), which makes an escarpment in the

west of Quilon Plateau (QP). This change in direction of isobaths may be of fault. Similar observation was made by *Yatheesh et al.* [2013] and *Nathaniel* [2013] and inferred as landward extension of the VFZ, whose expression continues southward into the Central Indian Basin. Recent studies by *Unnikrishnan et al.* [2017] has observed that VFZ is a crustal scale feature and plays a significant role on the sedimentation pattern on either side (east and west) of the fracture zone.

Furthermore, step like features, similar to Horst and Graben has also been identified in Zone II of profile C–C' of Figure 6.

4.3. Zone III

The southern part of the study area belongs to the Trivandrum block, which lies below the ACSZ. The continental shelf in this part narrows down and becomes 50 km at off Trivandrum. Moreover, Zone III is important in global perspective due to the gravitational geoidal anomaly. Three set of drainages occurring south of QP are parallel to sub parallel in nature, which meets its higher order streams in right angle having a trend of NW–SE. (Figure 5, profile D–D' of Figure 6). The southern part of QP has a gentle slope with step-like features/terraces. These terraces are part of Allepy Trivandrum Terrace Complexes (ATTC), [*Biswas, 1987; Yatheesh et al., 2006, 2013*]. It is inferred that these terraces are rifted and sheared segment of pre-drift plate boundary related to India–Madagascar separation during 86.5 Ma.

4.4. Satellite Derived Gravity Data

The lithological contrast is clearly visible in gravity data downloaded from satellite altimetry data (Figure 7). In Figure 7, ridge-like features identified in Zone I of study area similar to the bathymetric studies. Earlier studies imply that these structural features are deep seated faults with rifted fragment of Western continental margin of India formed during the north ward movement and during the Paleogene, these ridges exposed to sub-areal erosion create flat top mounts and in the Neogene it has been submerged, [*Gopala Rao et al., 2010; Rao and Wagle, 1997; Ramana et al., 2015*].

In Zone II we observed that the gravity changes

rapidly and it overlaps very well with early identified VFZ. Furthermore, gravity data clearly shows the concealed lithology and Escarpment in QP.

In Zone III, negative gravity anomaly has been observed; [*Rao et al., 2014*] described this negative geoidal anomaly towards south of Indian Continent as Indian Ocean Geoid Low. Moreover, we observed gravitational value in sigmoidal shape with the same value as in Trivandrum block in the south of India. So we are proposing that the sigmoidal portion in Zone III may be of the submerged part of continental crust of the Trivandrum block. This gravity data also proves the lithological contrast and possibility of fault/lineament which runs parallel to ACSZ and this observation is very well supported by [*Rao et al., 2017*].

4.5. Bathymetric Profiling From North to South and 3D Modelling

The bathymetric profiles from N–S exemplifies the major structural features described in Zones (I–III) for regional understanding of the west coast off Kerala (Figure 8 WCP-1 to 7). The Host and graben structures are magnified as step-like features in profiles. In Figure 9 and Figure 10 an attempt has been made to integrate data through vertical stacking of satellite derived GBM and Gravity data. It shows morphotectonic and neotectonic features of the west coast passive margin on a regional scale.

5. Conclusions

Transition from oceanic to continental crust is structurally complex and is often obscured by thick layers of sediment, with these sediment layers being of contrasting composition and density. To understand the morphotectonic setup of land and sea areas an integrated approach in analysis of geological, geophysical and spatial data can be used. Higher resolution satellite altimetry and gravity data along the west coast off Kerala has revealed many submarine morphotectonic features and its relation with the land. It serves as an important tool for the identification of neotectonic activities in and along the passive margin of the west coast off Kerala




Figure 7. Satellite derived Gravity map (from topex) with inferred faults and submerged continental crust/Trivandrum block?. The Sigmoidal shape of the submerged portion indicate major tectonic movement. The probable direction of movement of submerged block is also marked.

(Zone I, Zone II and Zone III). Other significant features like impact crater, submerged terraces etc. have also been observed using this technique. The morphotectonic evolution of drainage basins is a reliable indicator of tectonically reactivated strain zones, which helps to study submarine landslides and seismo-turbidites.

Acknowledgment. The authors wish to acknowledge Director General, Geological Survey of India (GSI) for his immense support. Further we are thankful to Deputy Director Generals and Directors of GSI for their encouragement. Also we are grateful to the reviewers and editors for their constructive comments for improving the manuscript.




Figure 8. DBM map along with profiles along the west coast off Kerala.




Figure 9. Three Dimensional 3D model of West Coast India with major tectonic features derived from GBM data.




Figure 10. An Integrated map of gravity and bathymetry(3D & 2D) along West Coast off Kerala.

References

- Ajayakumar, P., S. Rajendran, T. M. Mahadevan (2017), Geophysical lineaments of Western Ghats and adjoining coastal areas of central Kerala, southern India and their temporal development, *Geoscience Frontiers*, 8, No. 5, 1089–1104, [Crossref](#)
- Allen, P. A. (2008), From landscapes into geological history, *Nature*, 451, 274–276, [Crossref](#)
- Bastia, R., et al. (2010), Structural and tectonic interpretation of geophysical data along the eastern continental margin of India with special reference to the deep water petroliferous basins, *J. Asian Earth Sci.*, 39, 608–619, [Crossref](#)
- Belknap, W., R. J. Naiman (1998), A GIS and TIR Procedure to Detect and Map Wall-Base Channels in Western Washington, *J. Envi. Manag.*, 52, No. 2, 147–160, [Crossref](#)
- Bhattacharya, G. C., V. Yatheesh (2015), Plate-tectonic evolution of the deep ocean basins adjoining the western continental margin of India e a proposed model for the early opening scenario, *Mukherjee, S. (Ed.), Petrol. Geosci.: Indian Contexts, Springer Geology* [Crossref](#)
- Biswas, S. K. (1987), Regional tectonic framework, structure and evolution of the western marginal basins of India, *Tectonophysics*, 135, 307–327, [Crossref](#)
- Chatterjee, S., A. Goswami, C. R. Scotese (2013), The longest voyage: Tectonic, magmatic, and paleoclimatic evolution of the Indian plate during its northward flight from Gondwana to Asia, *Gond. Res.*, 23, 238–267, [Crossref](#)
- Chetty, T. R. K., T. Yellappa, M. Santosh (2016), Crustal architecture and tectonic evolution of the Cauvery Suture Zone, southern India, *Journal of Asian Earth Sciences*, 130, 166–191, [Crossref](#)
- Collins, A. S., C. Clark, D. Plavsa (2014), Peninsular India in Gondwana: the tectonothermal evolution of the Southern Granulite Terrane and its Gondwanan counterparts, *Gond. Res.*, 25, 190–203, [Crossref](#)
- Das, K. C. (2013), Mesozoic Enigma in Kerala-Konkan Basin: An alternate explanation for deep water sub-basalt reflections, *Proceeding of 10th Biennial International Conference & Exposition, 23–25 November 2013, Kochi, India* p. 352–359,
- Dhanya, V., G. Renoy (2015), Drainage development in Achankovil Shear Zone, South India, *Curr. Sci.*, 108, No. 6, 1151–1157.
- Duncan, R. A., M. Storey (1992), The life cycle of Indian Ocean hot spots, *Duncan, R. A. (Ed.), Synthesis of Results from Scientific Drilling in the Indian Ocean. American Geophys. Union Geophysical Monograph.*, 70 p. 91–103,
- Faruque, B. M., K. V. Ramachandran (2014), *The Continental Shelf of Western India*, 213–220 pp.,
- Ferraris, F., M. Firpo, F. J. Pazzaglia (2012), DEM Analyses and Morphotectonic Interpretation: The Plio-Quaternary Evolution of the Eastern Ligurian Alps, Italy, *Geomorphology*, 149–150, 27–40, [Crossref](#)
- Gahagan, L. M., et al. (1988), Tectonic fabric map of the ocean basins from satellite altimetry data, *Tectonophysics*, 155, 1–26, [Crossref](#)
- Girishbai, Drishya, Varghese Saju, A. C. Dinesh, Bhadraran Arun, R. K. Joshi (2017), Provenances of possible turbidites and volcanoclastic sediments in the northern part of Narcondam-Barren Basin, Andaman Sea, *Arab. J. Geosci.*, 10, 10–39, [Crossref](#)
- Girish, G., G. K. Ambili, N. P. Jesiya, L. Kuriachan (2016), Hydro-hypsometric analysis of tropical river basins south west coast of India using geospatial technology, *J. Mount. Science*, 15, No. 3, 939–946, [Crossref](#)
- Gopala Rao, D., A. L. Paropkari, K. S. Krishna, A. K. Chaubey, K. K. Ajay, V. N. Kodagali (2010), Bathymetric highs in the mid-slope region of the Western Continental Margin of India – structure and mode of origin, *Marine Geology*, 276, 58–70, [Crossref](#)
- Hashimi, N. H., et al. (1995), Holocene sea level curve and related climatic fluctuations for western Indian continental margin. An update, *Jour. Geol. Soc. India*, 46, 157–162. (Reprinted in “Vedic Saraswati: Evolutionary History of a Lost River of Northwestern India”, B. P. Radhakrishna and S. S. Merh (Eds.), Mem. Geol. Soc. India, 42, 297–302)
- Horton, R. E. (1945), Erosional development of streams and their drainage basins: hydrophysical approach to quantitative morphology, *Bull. Geo. Soc. Am.*, 56, 275–370, [Crossref](#)
- Imaki, H., T. Beechie (2008), *GIS Approaches for Channel Typing in the Columbia River Basin: Carrying Fine Resolution Data to a Large Geographic Extent American Geophys. Union, Abstract no H51A-0789*,
- Jensen, S. K., J. O. Dominique (1988), Extracting topographic structure from digital elevation data geographical information system analysis, *Photogrammetric Eng. & Remote Sensing*, 54, No. 11, 1593–1600.
- Karunakaran, C., T. M. Mahadevan (1971), Some aspects of the seismicity of peninsular shield, *Indian Jour. Power and River Valley Development, Sp. number Proc. Symp. Koyna Earthquake and Related Problems*, 1, 15–24.
- Kundu, S. N., D. S. Pattanaik (2011), Mapping on Land River channels up to the seafloor along the west coast of India, *Curr. Sci.*, 101, 958–961.
- Lin, W. T., W. C. Choub, C. Y. Lin, P. H. Huang, J. S. Tsai (2008), Win Basin: Using Improved Algorithms and the GIS Technique for Automated Watershed Modelling Analysis from Digital Elevation Models, *Int. J. Geograph. Infor. Sci.*, 22, No. 1, 47–69, [Crossref](#)
- Nathaniel, D. M. (2013), Hydrocarbon potential of sub-basalt Mesozoics of deepwater Kerala Basin, India, *Proceeding of 10th Biennial International Con-*

- ference & Exposition, 23–25 November 2013, Kochi, India* p. 1–7,
- Norton, I. O., J. G. Sclater (1979), A model for the evolution of the Indian Ocean and the break-up of Gondwanaland, *J. Geophys. Res.*, *84*, 6803–6830, [Crossref](#)
- O’Callaghan, J. F., D. M. Mark (1984), The extraction of drainage networks from digital elevation data, *Computer Vision, Graphics, and Image Processing*, *28*, No. 3, 323–344, [Crossref](#)
- Paull, C. K., W. I. Ussler, H. G. Greene, A. Keaten (2003), Caught in the act: The December 21 gravity flow event in Monterey Canyon, *Geo-Marine Letters*, *22*, 227–232, [Crossref](#)
- Pillay, K. R., et al. (2016), A possible impact crater from Outer Continental shelf off Cochin, Arabian Sea, India, *Curr. Sci.*, *111*, 1149–1151.
- Perna, Ramesh, et al. (2015), Approximation of Flow Patterns for Submarine Channel Systems in the Arabian Sea using a GIS Approach, Cloud Publications, *Int. J. Adv. Remote Sensing and GIS*, *4*, 1142–1160. (Article ID Tech-429 ISSN 2320-0243)
- Radhakrishna, T., et al. (1999), ⁴⁰Ar/³⁹Ar and K/Ar geochronology of the dykes from the south Indian granulite terrain, *Tectonophysics*, *304*, 109–129, [Crossref](#)
- Rahman, M. M., D. S. Arya, N. K. Goel (2010), Limitation of 90 m SRTM DEM in drainage network delineation using D8 method – a case study in flat terrain of Bangladesh, *Appl. Geomatics*, *2*, 49–58, [Crossref](#)
- Rajendran, C. P., K. Rajendran (1996), Low moderate seismicity in the vicinity of Palaghat Gap, South India and its implications, *Curr. Sci.*, *70*, 304–307.
- Ramana, M. V., Ana Desa Maria, T. Ramprasad (2015), Re-examination of geophysical data off Northwest India: Implications to the Late Cretaceous plate tectonics between India and Africa, *Mar. Geol.*, *365*, 36–51, [Crossref](#)
- Rao, V. P., B. G. Wagle (1997), Geomorphology and surficial geology of the western continental slope of India: a review, *Curr. Sci.*, *73*, 330–350.
- Rao, V. P., G. Rajagopalan, K. H. Vora, F. Almeida (2003), Late Quaternary sea level and environmental changes from relic carbonate deposits of the western margin of India, *Proc. Indian Acad. Sci. (Earth Planet. Sci.)*, *112*, No. 1, 1–25.
- Rao, B. P., M. R. Kumar (2014), Seismic evidence for slab graveyards atop the core–mantle boundary beneath the Indian Ocean geoid low, *Phys. Earth Planet. Inter.*, *236*, 52–59, [Crossref](#)
- Rao, P. N., R. M. Kumar, K. Arora (2017), Discovery of a massive ancient tectonic slab in the southeastern Indian Ocean: implications for the Indian Ocean geoid low, *Curr. Sci.*, *112*, 449–450.
- Reinfelds, I., T. Cohen, P. Batten, G. Brierley (2004), Assessment of Downstream Trends in Channel Gradient, Total and Specific Stream Power: A GIS Approach, *Geomorphology*, *60*, 403–416, [Crossref](#)
- Sandwell, D. T., W. H. F. Smith (1994), Bathymetric prediction from dense satellite altimetry and sparse shipboard bathymetry, *J. Geophys. Res.*, *99*, 803–824.
- Sandwell, D. T., W. H. F. Smith (1997), Global Sea Floor Topography from Satellite Altimetry and Ship Depth Soundings, *Science*, *277*, 1957–1962.
- Sandwell, D. T., W. H. F. Smith (2009), Global marine gravity from retracked Geosat and ERS-1 altimetry: Ridge segmentation versus spreading rate, *J. Geophys. Res.: Solid Earth*, *114*, No. 1, 1–18, [Crossref](#)
- Santosh, M., S. Maruyama, K. Sato (2009), Anatomy of a Cambrian suture in Gondwana: Pacific-type orogeny in southern India?, *Gond. Res.*, *16*, 321–341, [Crossref](#)
- Santosh, M., Q. Y. Yang, E. Shaji, M. Tsunogae, M. Ram Mohan, M. Satyanarayanan (2015), An exotic Mesoarchean microcontinent: the Coorg, Block, southern India, *Gond. Res.*, *29*, 105–135, [Crossref](#)
- Santosh, M., C. N. Hu, X. F. He, S. S. Li, M. Tsunogae, E. Shaji, G. Indu (2017), Neoproterozoic arc magmatism in the southern Madurai block, India: Subduction, relamination, continental outbuilding, and the growth of Gondwana, *Gond. Res.*, *45*, 1–42, [Crossref](#)
- Shareef, N. M., R. Mukhopadhyay, A. Sivasamy, et al. (2015), Delineation of buried channels of Bharathappuzha by single-channel shallow seismic survey, *Curr. Sci.*, *110*, No. 2, 153–156.
- Shreve, R. L. (1966), Statistical Law of Stream Numbers, *J. Geol.*, *74*, No. 1, 17–37, [Crossref](#)
- Singh, Y., B. John, G. P. Ganapathy, A. George, S. Harisanth, K. S. Divyalakshmi, S. Kesavan (2016), Geomorphic observations from southwestern terminus of palghat Gap, south India and their tectonic implications, *Journal of Earth System Science*, *125*, No. 4, 821–839, [Crossref](#)
- Strahler, A. N. (1952), Hypsometric (Area-Altitude) Analysis of Erosional Topology, *Geolog. Soc. of America Bulletin*, *63*, No. 11, 1117–1142.
- Subrahmanyam, C., S. Chand (2006), Evolution of the passive continental margins of India-A geophysical appraisal, *Gond. Res.*, *10*, 167–178, [Crossref](#)
- Subrahmanyam, V., M. V. Ramana, D. Gopala Rao (1993), Reactivation of Precambrian faults on the southwestern continental margin of India, evidence from gravity anomalies, *Tectonophysics*, *219*, 327–339, [Crossref](#)
- Sukumaran, P. V., et al. (2010), Marine sand in the south-west continental shelf of India, *Indian J. GeoMarine Sciences*, *39*, No. 4, 572–578.
- Umnikrishnan, P., M. Radhakrishna, G. K. Prasad (2017), Crustal structure and sedimentation history over the Alleppey platform, southwest continental margin of India: Constraints from multichannel seismic and gravity data, *Geoscience Frontiers*, *9*, 549–558, [Crossref](#)
- Vieux, B. E. (2001), Distributed Hydrologic Modeling Using GIS, Water Sci. and Tech. Library, *Dis-*

- tributed Hydrologic Modeling Using GIS. Water Science and Technology Library* 38, p. 1–17, **Crossref**
- Vijith, H., V. Prasannakumar, M. V. Ninu Krishna, P. Pratheesh (2015), Morphotectonics of a small river basin in the South Indian granulite terrain: An assessment through spatially derived geomorphic indices, *Georisk, Taylor & Francis*, 9, 187–199, **Crossref**
- Xu, J. P. (2010), Normalized velocity profiles of field-measured turbidity currents, *Geology*, 38, 563–566, **Crossref**
- Yatheesh, V., G. C. Bhattacharya, K. Mahender (2006), The terrace like features in the mid continental slope region off Trivandrum and a plausible model for India–Madagascar juxtaposition in immediate pre-drift scenario, *Gond. Res.*, 10, 179–185.
- Yatheesh, V., P. John Kurian, G. C. Bhattacharya, S. Rajan (2013), Morphotectonic architecture of an India–Madagascar breakup related anomalous submarine terrace complex on the southwest continental margin of India, *Mar. Pet. Geol.*, 46, 304–318, **Crossref**
-
- Arun Bhadrans, Geodata Division, Central Head Quarters, Geological Survey of India, Kolkata, India
- Drishya Girishbai, International Affairs & IGC, Central Head Quarters, Geological Survey of India, Kolkata, India. (geodrish@gmail.com)